

Independent losses and duplications of autophagy-related genes in fungal tree of life

Qinhu Wang,¹ Huiquan Liu,¹ Huaijian Xu,¹
Ruonan Hei,¹ Shijie Zhang,² Cong Jiang¹ and
Jin-Rong Xu^{1,3*}

¹State Key Laboratory of Crop Stress Biology for Arid Areas and College of Plant Protection, Northwest A&F University, Yangling, Shaanxi, 712100, China.

²School of Life Sciences, Zhengzhou University, Zhengzhou, Henan, 450001, China.

³Department of Botany and Plant Pathology, Purdue University, West Lafayette, IN, 47907, USA.

Summary

Autophagy is important for growth, development and pathogenesis in fungi. Although autophagic process is generally considered to be conserved, the conservation and evolution of ATG genes at kingdom-wide remains to be conducted. Here we systematically identified 41 known ATG genes in 331 species and analyzed their distribution across the fungal kingdom. In general, only 20 ATG genes are highly conserved, including most but not all the yeast core-autophagy-machinery genes. Four functional protein groups involved in autophagosome formation had conserved and non-conserved components, suggesting plasticity in autophagosome formation in fungi. All or majority of the key ATG genes were lost in several fungal groups with unique lifestyles and niches, such as Microsporidia, *Pneumocystis* and *Malassezia*. Moreover, majority of ATG genes had A-to-I RNA editing during sexual reproduction in two ascomycetes and deletion of *FgATG11*, the ATG gene with the most editing sites in *Fusarium* affected ascospore releasing. Duplication and divergence also was observed to several core ATG genes, such as highly divergent *ATG8* paralogs in dermatophytes and multiple *ATG15* duplications in mushrooms. Taken together, independent losses and duplications of ATG genes have occurred throughout the fungal kingdom and variations in autophagy exist among different lineages and possibly different developmental stages.

Received 8 June, 2018; accepted 16 October, 2018. *For correspondence. E-mail jinrong@purdue.edu; Tel. +1-765-496-6918; Fax +1-765-494-5896.

Introduction

Autophagy is an intracellular degradation process that is critical for cellular homeostasis (Farre and Subramani, 2016). It is a nutrient recycling mechanism for dealing with the superfluous and damaged organelles (Feng *et al.*, 2014). The autophagy process consists of three sequential steps: cargo sequestration, degradation and utilization of degradation products (Mizushima, 2007). For cargo sequestration, the pre-autophagosomal structure (PAS) is activated to generate the autophagosome with double membranes. PAS is very close to the vacuolar membrane (Yoshimori and Noda, 2008). In the degradation step, autophagosomes fuse with lysosomes or vacuoles and form the degrading structure known as autophagolysosomes for cargo degradation (Eskelinen and Saftig, 2009). In the last step, the nutrient monomeric units deriving from the degradation of macromolecules are exported to the cytosol for recycling (Mizushima, 2007). According to the cargoes and degradation manner, autophagy is divided into three primary types: macroautophagy, microautophagy and chaperone-mediated autophagy (Mizushima *et al.*, 2008). The macroautophagy is primarily a non-selective process and commonly referred to as autophagy.

The autophagic machinery consists of the autophagy-related (ATG) genes (Feng *et al.*, 2014). As a model for autophagy studies, many groundbreaking and fundamental knowledge of the autophagic processes have been characterized in the budding yeast *Saccharomyces cerevisiae*, including the identification of the ATG genes such as *ATG1* and *ATG8* (Feng *et al.*, 2014). *ATG1* is the first ATG gene that was shown to be essential for the accumulation of autophagic bodies and autophagy in yeast (Tsukada and Ohsumi, 1993). It encodes a protein kinase that is involved in the recruitment and release of other autophagy-related proteins from the PAS (Yoshimori and Noda, 2008). *ATG8* encodes an ubiquitin-like protein that is essential for autophagosome formation and selective cargo recruitment. The GFP-Atg8 fusion is commonly used as a marker to follow autophagy (Xie *et al.*, 2008; Shpilka *et al.*, 2011).

Autophagy has a variety of functions in eukaryotes. In mammalian cells, autophagy is known to be involved in

embryonic development, cell differentiation, immunity and inflammation (Mizushima and Levine, 2010; Levine *et al.*, 2011; Kanayama and Shinohara, 2016). In plants, autophagy is important for seedling establishment, plant development, senescence, stress resistance, metabolism and reproduction (Liu *et al.*, 2005; Bassham *et al.*, 2006; Avila-Ospina *et al.*, 2014; Michaeli *et al.*, 2016). In fungi, autophagy affects growth, morphology, development, lipid turnover and sexual reproduction (Kershaw and Talbot, 2009; Pollack *et al.*, 2009; Nguyen *et al.*, 2011; Josefsen *et al.*, 2012; Liu *et al.*, 2012; Duan *et al.*, 2013; Voigt and Poggeler, 2013; Lv *et al.*, 2017). In fungal pathogens, autophagy is also known to be important for plant or animal infection. For example, deletion any of the 16 ATG genes required for nonselective autophagy in the rice blast fungus *Magnaporthe oryzae* affected its virulence (Kershaw and Talbot, 2009). Mutants of many ATG genes were defective in toxin production and plant infection (Nguyen *et al.*, 2011; Wang *et al.*, 2011; Josefsen *et al.*, 2012; Lv *et al.*, 2017) in another ascomycete *Fusarium graminearum*, the main causal agent of Fusarium head blight (FHB). In human pathogen *Cryptococcus neoformans*, knockdown of ATG8 results in attenuated virulence in a mouse model of infection (Hu *et al.*, 2008). In the corn smut fungus *Ustilago maydis*, ATG1 and ATG8 are required for the full virulence and complete symptom development (Nadal and Gold, 2010).

Although many ATG genes are conserved from yeast to humans (Meijer *et al.*, 2007), several of them such as ATG25 are yeast specific (Monastyrska *et al.*, 2005). Furthermore, during the study of kinome in *F. graminearum* (Wang *et al.*, 2011), we found that the Atg1 kinase, an essential regulator of autophagy, is not present in some of the fungal genomes. Considering the fact that major knowledge of autophagy is based on yeast studies and the importance of autophagy in fungal development and pathogenesis, it is important to investigate the conservation and distribution of ATG genes across the fungal kingdom to better understand the evolution and regulation of autophagy in fungi. In this study, we systematically identified and analyzed 41 known ATG genes in 331 fungal species. The distribution of each ATG gene was analyzed across the fungal kingdom. Massive and specific losses of ATG genes were observed in fungi belonging to different phyla. Duplication of several ATG genes, including ATG8, ATG15, ATG18, ATG20 and ATG22, also were observed in different fungal groups. In Arthrodermataceae, two Atg8 homologues are likely derived from an ancient duplication event because of their phylogenetic relationship. We also found that majority of the ATG genes had multiple A-to-I RNA editing sites during sexual reproduction in two filamentous ascomycetes. Deletion of ATG11, the ATG gene with the most RNA editing sites in

Fusarium graminearum, had no effect on growth, infection, conidiation and ascus development but resulted in defects in ascospore releasing. Taken together, our results showed that independent losses and duplication of ATG genes have occurred in fungal tree of life and some unrelated fungi with unique growth niches and reduced genome sizes lack any or key ATG genes and autophagy. Furthermore, post-transcriptional modifications by RNA editing may affect stage-specific functions of ATG genes and autophagy during sexual reproduction in fungi.

Results

The distribution of ATG genes varies significantly among different fungal classes

To comprehensively analyze the ATG genes in kingdom Fungi, we search for homologues of 36 *S. cerevisiae* ATG genes (www.yeastgenome.org) that are categorized into core machinery, selective autophagy specific and nonselective autophagy specific group (Lynch-Day and Klionsky, 2010), and homologues of ATG25, ATG28, ATG30, ATG35 and ATG37 of *Hansenula polymorpha* or *Pichia pastoris* (Monastyrska *et al.*, 2005; Meijer *et al.*, 2007) (Supporting Information Table S1). Their orthologs in the genomes of 331 sequenced fungal species, including 227 Ascomycota, 72 Basidiomycota and 32 lower fungi (Fig. 1A; Supporting Information Table S2) were identified.

Based on their conservation (Fig. 1B) and distribution (Fig. 1C) across the kingdom Fungi, these ATG genes were categorized into three groups. The first group contains 20 highly conserved ATG genes, including ATG1-9, ATG11-13, ATG15, ATG18, ATG20-22, ATG24 and ATG26-27 (Fig. 1B and C). Their orthologs are present in over 80% of the fungal species that have been sequenced. Whereas most of them (14 out of 20) are core machinery ATG genes, six of them are selective autophagy-specific genes. The second group includes seven ATG genes, ATG10, ATG16, ATG17, ATG28, ATG29, ATG33 and ATG37 that are present in Ascomycota but not in Basidiomycota or basal fungal groups (Fig. 1B and C). Group III consists of 14 ATG genes that are present only in Saccharomycetes species (Fig. 1B and C), including ATG14, ATG38 and ATG41 that are core autophagy genes in *S. cerevisiae*.

Interestingly, examination of the 46 sequenced ascomycetous yeast species revealed that some of the ATG genes are not well conserved in Saccharomycetes. Among the highly conserved core machinery ATG genes, ATG22 that encodes a vacuolar membrane protein is lost in the Debaryomycetaceae family (Supporting Information Fig. S1). The seven group II ATG genes, including two yeast core




Fig. 1. Autophagy-related genes in different classes of fungi.

A. The number of fungi belonging to different fungal phyla surveyed in this study.

B. The number of fungal species containing 0 (grey), 1 (blue) or at least 2 copies (pink) of individual ATG genes (*ATG1*–*ATG41* labelled as 1–41 at the bottom). The numbers of highly conserved, conserved and Saccharomycetes-specific ATG genes were shaded in red, green and yellow respectively. These ATG genes also were labelled on the top with C for core machinery, S for selective autophagy specific and N for nonselective autophagy specific genes based on the classification of their orthologs in the budding yeast.

C. The conservation of ATG genes in different lineages across the kingdom Fungi. The dendrogram on the left shows the phylogenetic relationship of the 331 fungi belonging to labelled fungal classes or phyla. The numbers of species in each class or phylum are indicated in the bracket. Individual pie charts in the matrix show the ratio of fungal species in each fungal groups with (blue) versus without (grey) individual ATG genes. [Color figure can be viewed at wileyonlinelibrary.com]

machinery ATG genes, *ATG10* and *ATG16*, were absent in 17–32 Saccharomycetales species (Supporting Information Fig. S1). Taken together, although the core machinery ATG genes are generally conserved, different fungal classes differ in the conservation of selective or non-selective autophagy-specific genes.

Plasticity in the components of protein complexes involved in autophagosome formation

The formation of autophagosomes involves four complexes of Atg proteins and two ubiquitin-like (Ubl) conjugation systems (Suzuki *et al.*, 2016). Except the

Atg2–Atg8 complex, all the other functional groups consist of components that are not well conserved in fungi (Fig. 2). For the Atg1 complex, components of the Atg1–Atg13 regulatory subunit are highly conserved across the fungal kingdom. However, components of the Atg17–Atg29–Atg31 scaffolding sub-complex are only conserved in ascomycetes. For the autophagy-specific phosphatidylinositol (PI) 3-kinase (PI3K) complex, Atg14 and Atg38, are specific for Saccharomycetes (Fig. 2). Similarly, Atg41 of the Atg9 vesicle complex is present only in Saccharomycetes. For the two Ubl conjugation systems required for the formation of autophagosomes (Ohsumi, 2001; Geng and Klionsky, 2008), the E2-like enzyme




Fig. 2. Conservation analysis of important complex or process involved in autophagosome formation. For each protein complex involved in different processes of autophagosome formation as diagrammed underneath, the Atg proteins highlighted in red, green or yellow represent the ATG genes that are highly conserved, conserved or *Saccharomyces*-specific respectively. For the PI3K complex I, the components shaded in grey (Vps15 and Vps34) were not covered in this analysis. [Color figure can be viewed at wileyonlinelibrary.com]

Atg3 is highly conserved across the fungal kingdom in the Atg8-conjugating system. However, the E2-like enzyme Atg10 of the Atg12-conjugating system is only present in some of the Ascomycota species (Fig. 2; Supporting Information Table S2). Therefore, despite their conserved roles in autophagosome formation, these Atg protein complexes or systems have significant variations in compositions. Overall, among the 20 ATG genes involved in the autophagosome formation in *S. cerevisiae* (Mizushima, 2007; Yao *et al.*, 2015; Suzuki *et al.*, 2016), 12 of them are highly conserved in fungi. The other eight are either only present in Ascomycota or specific for *Saccharomyces* (Fig. 2), suggesting that autophagosome formation and its regulation vary among different fungal groups.

The ATG genes are completely or almost completely lost in Microsporidia

Microsporidia is a phylum of spore-forming unicellular fungi that infect human and other animals (Keeling, 2009). All the Microsporidia species are obligate, intracellular pathogens. Among the 20 Microsporidia species analyzed, 11 of them have only the ATG15 gene (Fig. 3). Six species, including *Nosema apis* and *Edhazardia aedis*, lack any ATG gene (Fig. 3). Interestingly, *Mitosporidium daphnia* that still possesses mitochondria (Haag *et al.*, 2014) has six ATG genes (ATG1/2/7/8/9/15), which is the most among the Microsporidia species (Fig. 3). Intracellular obligate pathogens depending on host cells for growth and reproduction often have a reduced genome size (Casadevall, 2008; Corradi, 2015; Ma *et al.*, 2016). It is likely that Microsporidia may no longer need autophagy and have lost most or all the ATG genes. Furthermore, differences among Microsporidia

species in the number of ATG genes suggest that independent gene loss events likely occurred to some of them during evolution.

Most of the core machinery ATG genes are lost in Pneumocystis species

Like Microsporidia, *Pneumocystis* species are obligate, intracellular pathogens. As a genus in Taphrinomycotina, *Pneumocystis* consists of species that infect lungs of human and other mammals (Ma *et al.*, 2016). About 33 out of the 41 ATG genes were absent in all three *Pneumocystis* species analyzed, including 12 of the 20 highly conserved group I fungal ATG genes (Fig. 3). Only three ATG genes, ATG15, ATG20 and ATG22, are commonly present in *P. jirovecii*, *P. murina* and *P. carinii*. However, these three *Pneumocystis* species differ in the presence of ATG6, ATG18, ATG21, ATG24 and ATG27 (Fig. 3). These results indicate that whereas most of the autophagy-related genes might be lost in their common ancestor, *Pneumocystis* also had species-specific gene loss events for some of the ATG genes.

Massive losses of ATG genes also occurred in Malassezia species

To our surprise, massive losses of ATG genes also were observed in all the three *Malassezia* species that have been sequenced (Fig. 3). As a genus in the Malasseziomycetes of Ustilagomycotina, *Malassezia* consists of free-living basidiomycete yeasts that are associated with animal skin disorders (Ashbee and Evans, 2002). *M. sympodialis*, an atopic eczema-associated pathogen, has only 7 ATG genes (Fig. 3). The other two *Malassezia* species have lost fewer ATG genes

than *M. sympodialis*. *M. globosa*, a dandruff-causing fungus and *M. pachydermatis*, a fungus frequently found on the skin and in the mucosa and ear canals of dogs, have 12 and 13 ATG genes respectively (Fig. 3). It is likely that these *Malassezia* species are defective in autophagy due to the massive losses of many highly conserved ATG genes.

ATG1 and 8 other highly conserved ATG genes are absent in a nematode trapping fungus

ATG1 encodes a protein kinase that is essential for the initiation of autophagosomes and formation of cytoplasm-to-vacuole targeting vesicles (Cvt, a selective autophagy). Interestingly, *Drechslerella stenobrocha* (Orbiliomycetes) appears to lack a distinct *ATG1* ortholog (Supporting Information Table S2). Bidirectional blastp and tblastn searches also failed to identify *ATG1* gene in its genome. *D. stenobrocha* also lacks 8 other highly conserved ATG genes, including *ATG4*, *ATG5*, *ATG9*, *ATG11*, *ATG15*, *ATG18*, *ATG20* and *ATG21*. Although it still has 16 ATG genes, the loss of *ATG1* and 8 other highly conserved ATG genes likely eliminated autophagy in this fungus. *D. stenobrocha* is a nematode trapping

fungus that forms constricting rings (Liu et al., 2014). It favours environments rich in organic compounds and its genome is 11 Mb smaller than that of *Arthrobotrys oligospora*, a close-related nematode trapping fungus (Liu et al., 2014). Loss of *ATG* genes in *D. stenobrocha* may be related to its adaption to the organic compound-rich environments and reduction in the genome size.

Many of the ATG genes are duplicated in Mucoromycotina fungi

Mucoromycotina contains several species that have been used for the production of fermented soy food in Asia and Africa (Dolatabadi et al., 2016). Some Mucoromycotina species are emerging life-threatening human pathogens (Roden et al., 2005). Seven highly conserved group I fungal ATG genes (*ATG8*, *ATG13*, *ATG15*, *ATG18*, *ATG20*, *ATG21* and *ATG24*) were duplicated in Mucoromycotina species (Fig. 3), which may be related to the ancient genome duplication events in Mucoromycotina (Schwartz et al., 2014). Interestingly, some of these duplicated ATG genes have more than two copies. For example, *ATG24* have three copies in almost all the Mucoromycotina species analyzed, indicating that




Fig. 3. Massive losses and duplications of ATG genes in specific fungal groups. The branches of Microsporidia, *Pneumocystis*, *Malassezia* and Mucoromycotina are in red, green, purple and blue, respectively, in the dendrogram showing their phylogenetic relationship on the left. The copy numbers of individual ATG genes (ATG1–ATG41, labelled on the top) in each species vary from 0 to 4. *Fusarium graminearum*, *Magnaporthe oryzae*, *Neurospora crassa*, *Komagataela pastoris*, *Ogataea parapolymorpha* and *Saccharomyces cerevisiae* are included for comparison. [Color figure can be viewed at wileyonlinelibrary.com]

duplications of *ATG24* likely occurred twice before the divergence of these species. Therefore, both genome-wide and gene-specific duplications events may have contributed to the expansion of these *ATG* genes in Mucoromycotina.

Ancient duplication and divergence of ATG8 in dermatophytes

Atg8 is a ubiquitin-like protein that is essential for the generation of double-membrane autophagosomes (Shpilka *et al.*, 2011). The *ATG8* orthologs are present in all the fungal genomes analyzed but Microsporidia and *Pneumocystis* species. Notably, several dermatophyte fungi in

Arthrodermataceae, including *Trichophyton rubrum*, one of the causal agent of athlete's foot, have two *ATG8* genes (Fig. 4A). Sequence alignment and phylogenetic analysis showed that Atg8a is highly similar to other fungal Atg8 proteins. In contrast, Atg8b has a high sequence divergence with typical fungal Atg8 proteins and appears to be unique to these dermatophytes (Fig. 4A and B). Although their amino acid sequences are not highly similar, both Atg8a and Atg8b have an ubiquitin-like protein folding (Fig. 4C). Furthermore, both *ATG8a* and *ATG8b* were expressed in *T. rubrum* based on the published RNA-seq data (Xu *et al.*, 2015) although the later had a much lower expression level (Fig. 4D). It is possible that both *ATG8* alleles encode ubiquitin-like proteins and have different or overlapping functions in Arthrodermataceae species.




Fig. 4. The divergence of *ATG8* genes in dermatophytes.

A. Sequence divergence and phylogenetic relationship between the two Atg8 proteins, Atg8a and Atg8b, in dermatophytes. The phylogenetic tree of fungal Atg8 proteins was inferred by PhyML 3.1 and the SH-like support values of branches, from lower and higher, were designated with circles from green to red. The abbreviations for the species used in the phylogenetic analysis are listed in Supporting Information Table S3. Atg8b proteins are specific for Arthrodermataceae and form a distinct clade.

B. Sequence alignment of Atg8a and Atg8b from six dermatophytes and Atg8 orthologs from *F. graminearum* (Fg), *M. oryzae* (Mo) and *N. crassa* (Nc).

C. The predicted 3D structures of *T. rubrum* Atg8a and Atg8b.

D. IGV sashimi plots showing the coverage of RNA-seq reads for *ATG8a* and *ATG8b* in *T. rubrum*. The density of RNA-seq reads are shown in red, the gene models are shown in blue, and the junction reads are shown as arcs with the read number. [Color figure can be viewed at wileyonlinelibrary.com]

ATG20 is duplicated in the fission and budding yeasts but not in filamentous fungi

In *S. cerevisiae*, *ATG20* (*SNX42*) and *SNX41* are two members of the sorting nexin family. Homologues of these two genes are also present in most of the Saccharomycetes species except the members of the Dipodascaceae family, such as *Geotrichum candidum* and *Yarrowia lipolytica* (Supporting Information Fig. S1; Supporting Information Table S2). Phylogenetic analysis revealed that *ATG20* and *SNX41* are two paralogs

derived from a duplication occurred in the common ancestor of Saccharomycetaceae after its divergence from Dipodascaceae (Fig. 5A).

Most filamentous ascomycetes and basidiomycetes had only one *ATG20*. However, the Schizosaccharomycetes species also have two *ATG20* homologues (Supporting Information Fig. S1; Supporting Information Table S2). Phylogenetic analysis showed that the two *ATG20* paralogs in *S. cerevisiae* and *S. pombe* were derived from independent gene duplication events (Fig. 5A). The two paralogs in




Fig. 5. Independent duplication of *ATG20* and *ATG18* in Saccharomycetes and Schizosaccharomycetes.

A. Phylogenetic tree of yeast *Atg20* and *Snx41* (two paralogs) and their orthologs from other ascomycetes and basidiomycetes. Except *Geotrichum candidum* (Gc) and *Yarrowia lipolytica* (Yl) that has a single copy, the other Saccharomycetes species have both *Atg20* and *Snx41* orthologs. *S. pombe* and three other Schizosaccharomycetes species also have the *Atg20* and *Snx41* paralogs. However, the paralogous *Atg20* and *Snx41* in Saccharomycetes and Schizosaccharomycetes are in different clades, suggesting their independent origins. Furthermore, both *Atg20* and *Snx41* clades are not clustered or aligned with the single copy of *Atg20/Snx41* in their closely related species such as *Y. lipolytica* or *T. deformans*, suggesting the occurrence of sequence divergence to both *Atg20* and *Snx41* in these yeast species.

B. Phylogeny of *S. cerevisiae* *Atg18* and *Atg21* and their orthologs from Saccharomycetes species.

C. Phylogeny of *S. pombe* *Atg18* proteins and their orthologs from other Schizosaccharomycetes species. The *Atg18/21* orthologs from *Saitoella complicata* (Sco) and *Taphrina deformans* (Td) are included for comparison with other Taphrinomycotina species. The phylogenetic tree was inferred by PhyML 3.1, and the SH-like support values of branches, from lower and higher, were designated with circles from green to red. The abbreviations for the species used in the phylogenetic analysis are listed in Supporting Information Table S3. [Color figure can be viewed at wileyonlinelibrary.com]

Schizosaccharomycetes species are likely derived from a duplication event occurred in their common ancestor after its divergence from other Taphrinomycota lineages. In *S. cerevisiae*, whereas Atg20 is required for both pexophagy and pexophagy-independent endosomal retrieval trafficking, Snx41 is only involved in the latter (Hettema *et al.*, 2003; Deng *et al.*, 2012). The two *ATG20* paralogs also differ in their functions in *S. pombe* (Zhao *et al.*, 2016). *MoATG20*, the only *ATG20* ortholog in *M. oryzae*, was shown to play roles in both protein sorting (Snx41 function) and pexophagy (Atg20 function) (Deng *et al.*, 2013). The dual functions of Atg20 proteins are likely conserved in other filamentous fungi with a single *ATG20* homologue.

ATG18 is also duplicated in the ascomycetous yeasts but not in filamentous fungi

Atg18 and Atg21 are two paralogous proteins with WD-40 repeats in *S. cerevisiae* (Krick *et al.*, 2008; Nair *et al.*, 2010). Whereas Atg18 is involved in both autophagy and Cvt pathways, Atg21 is only required for the latter (Barth *et al.*, 2001; Stromhaug *et al.*, 2004). Paralogous Atg18 and Atg21 also are present in most Saccharomycetes species except *Y. lipolytica* (Supporting Information Fig. S1; Supporting Information Table S2). Phylogenetic analysis showed that *ATG18* and *ATG21* also were from a duplication in the last common ancestor of Saccharomycetaceae (Fig. 5B). Most filamentous fungi have a single *ATG18* ortholog. Interestingly, *S. pombe* and other Schizosaccharomycetes have three genes that are homologous to *ATG18* or *ATG21* (Supporting Information Table S2). Phylogenetic analysis showed that these three *ATG18/21* homologues likely are derived from two separate duplication events occurred in the common ancestor of Schizosaccharomycetes (Fig. 5C). The duplications of these *ATG* genes in unicellular yeasts but not in multicellular filamentous fungi may be related to their differences in life styles or vegetative growth and development.

Lineage-specific gene duplications of two ATG genes required for autophagic body breakdown

Two *ATG* genes, *ATG15* and *ATG22*, are involved in the breakdown and release of autophagic bodies (Lynch-Day and Klionsky, 2010). Our analysis showed that some Chytridiomycota species have one but others have two *ATG15*. Whereas majority of Ascomycetes have only one, most Agaricomycetes have three *ATG15* homologues, which are likely derived from two separate, sequential gene duplication events occurred in the common ancestor of Agaricomycetes fungi (Fig. 6A). In addition to these three ancient paralogs, some Agaricomycetes species also have recent duplications of *ATG15* (Fig. 6A). For example, *Pisolithus microcarpus* (Pm), an ectomycorrhizal fungus, has

five *ATG15* genes, including three ancient alleles and two alleles derived from recent gene duplication events.

In contrast to *ATG15*, most Agaricomycotina and Pucciniomycotina fungi have only one but many Ascomycetes have two *ATG22* genes. It appears that *ATG22* was duplicated in the ancestor of filamentous ascomycetes, resulting in two ancestral paralogs, *ATG22a* and *ATG22b* (Fig. 6B). Whereas *ATG22a* is well-conserved, *ATG22b* was lost in most branches of Pezizomycotina, including all the Sordariomycetes species (Fig. 6B). Interestingly, two copies of *ATG22* are present in *Puccinia* rust fungi (Fig. 6B) but none of the smut fungi in Ustilaginomycotina has *ATG22* (Supporting Information Table S2). These data suggest that filamentous or higher fungi may have different approaches in the releasing of recycled nutrients by lineage-specific duplication of *ATG15* or *ATG22*.

Most ATG genes are subjected to RNA editing during sexual reproduction in Fusarium graminearum and Neurospora crassa

Autophagy is known to be important for sexual reproduction in fungi (Wang *et al.*, 2011; Voigt and Poggeler, 2013; Lv *et al.*, 2017). Recently, it has been reported that A-to-I RNA editing occurs specifically during sexual reproduction in *F. graminearum*, *N. crassa* and other filamentous ascomycetes (Liu *et al.*, 2016, 2017). The genes involved in autophagy were significantly enriched among the genes with nonsynonymous editing events in these two fungi, suggesting that A-to-I RNA editing likely plays a stage-specific role in autophagy during sexual reproduction. When the published strand-specific RNA-seq data were analyzed, transcripts of most *ATG* genes had A-to-I RNA editing events in both *F. graminearum* and *N. crassa* (Fig. 7A). In *F. graminearum*, among the 24 *ATG* genes with A-to-I RNA editing, 22 of them had nonsynonymous editing resulting in amino acid changes. In *N. crassa*, all the 23 *ATG* genes with A-to-I RNA editing had nonsynonymous editing events (Fig. 7A). Many of these *ATG* genes had multiple editing events that specifically occurred during sexual reproduction, such as 27 editing sites in *FgATG11* and 31 editing sites in *NcATG2*. Interestingly, three *ATG* genes in *F. graminearum* (*FgATG7*, *FgATG15* and *FgATG28*) and six *ATG* genes in *N. crassa* (including 5 selective autophagy genes) had stop-codon-loss editing events, which will cause read through and possibly produce longer proteins than normal ones specifically during sexual reproduction (Fig. 7B).

In comparison between *F. graminearum* and *N. crassa*, a total of 10 editing events identified in 9 *ATG* genes were found to be conserved, including a stop-loss editing site in *ATG28* (Fig. 7). Two of these conserved editing sites in *ATG26* and *ATG13* resulted in the recoding of serine to glycine, which may affect the phosphorylation or structure of these Atg proteins. Interestingly, a conserved




Fig. 6. Phylogenetic analysis of Atg15 and Atg22 proteins involved in autophagic body breakdown in filamentous fungi.

A. Phylogeny of Atg15 homologues from different ascomycetes and basidiomycetes. Many Agaricomycetes have three Atg15 paralogs.

B. Phylogeny of Atg22 homologues for representative ascomycetes and basidiomycetes to show the duplication of Atg22 in Pezizomycotina and *Puccinia* species. The phylogenetic tree was inferred by PhyML 3.1, and the SH-like support values of branches, from lower and higher, were designated with circles from green to red. The abbreviations for the species used in the phylogenetic analysis are listed in Supporting Information Table S3. [Color figure can be viewed at wileyonlinelibrary.com]

editing event resulting in the isoleucine to valine (I32V) recoding was identified in *ATG8*, leading a similar consequence to the difference between Atg8a and Atg8b in dermatophytic fungi. Taken together, these data suggest that RNA editing may affect autophagy during sexual reproduction in both *F. graminearum*, *N. crassa* and other filamentous ascomycetes.

Mutations affecting the I32V and I41V editing events in FgATG8 had no obvious effects on its function during sexual reproduction

To determine the importance of two editing events at A94 and A121 (resulting in the I32V and I41V nonsynonymous

changes), we first generated the non-editable alleles of *FgATG8* (FGSG_10740) by introducing the T93 to G and T120 to C mutations. Because of the preference of U at the -1 position for A-to-I editing (Liu *et al.*, 2016), the T93G and T120C mutations likely will eliminate the editing of A94 and A121 in *FgATG8* transcripts. The resulting *FgATG8*^{G93} and *FgATG8*^{C120} alleles were transformed into the *Fgatg8* mutant generated by the split-marker approach (Catlett *et al.*, 2002; Josefsen *et al.*, 2012). When assayed for sexual reproduction, whereas the *Fgatg8* mutant was sterile, both the *Fgatg8*/*FgATG8*^{G93} and *Fgatg8*/*FgATG8*^{C120} transformants were similar to the wild type in perithecial formation and ascospore development (Supporting Information Fig. S2). We then




Fig. 7. The A-to-I RNA editing sites in ATG genes in *F. graminearum* and *N. crassa*. The RNA editing sites identified in transcripts of marked ATG genes during sexual reproduction in *F. graminearum* and *N. crassa*. The number of RNA editing sites in each ATG gene is marked in the bracket. The red and blue vertical lines represent the nonsynonymous and synonymous editing sites respectively. Black arrows mark the editing sites conserved between *F. graminearum* and *N. crassa*. [Color figure can be viewed at wileyonlinelibrary.com]

introduced the A94 to G and A121 to G mutations to *FgATG8* to generate the edited *FgATG8*^{G94} and *FgATG8*^{G121} alleles and transformed them into the *Fgatg8* mutant. The resulting *Fgatg8/FgATG8*^{G94} and *Fgatg8/FgATG8*^{G121} transformants also were normal in sexual reproduction (Supporting Information Fig. S2). These results indicated that mutations affecting the two A-to-I editing sites in *FgATG8* individually had no obvious effects on its functions during sexual reproduction. Because the editing levels at A94 (10.3%) and A121 (13.0%) in *FgATG8* were relatively low, it is not surprising that mutations affecting these two editing sites had no obvious effects under laboratory conditions. However, nonsynonymous editing events with low editing levels tend to confer to protein heterozygosity advantage in fungi (Liu *et al.*, 2017). Therefore, it remains possible that these two editing

events are advantageous for *FgAtg8* functions during sexual reproduction under field conditions.

Deletion of the FgATG11 results in sexual-specific defects in F. graminearum

Transcripts of *FgATG11* (FGSG_15734) had 27 A-to-I RNA editing sites, the most among all the edited ATG genes in *F. graminearum*. Two of these editing sites are conserved in *NcATG11* of *N. crassa* (Fig. 7). These data suggested that editing of this selective autophagy specific gene may be important for its function during sexual reproduction. To test this hypothesis, we generated the mutants deleted of *FgATG11* by the split marker approach (Catlett *et al.*, 2002). Four *Fgatg11* deletion mutants were





Fig. 8. Phenotypes of the *Fgatg11* mutant in *F. graminearum*. The wild-type strain PH-1 and *Fgatg11* deletion mutant of *F. graminearum* were assayed for colony growth (A, three-day-old cultures), virulence on wheat heads (B, sampled 14-days post-inoculation), disease index (C, diseased spikelets per wheat heads), DON production (D), conidiation (E), the germination rate of conidia (F), perithecia development on carrot agar plates (G), asci and ascospores (H, bar = 20 μ M), the germination rate of ascospores (I) and ascospore discharge from perithecia at 10 days post-fertilization (J). The defect in ascospore discharge was detected in all the four *Fgatg11* mutant strains repeatedly. In C, D, E, F and I, error bars represent the standard deviations from at least 3 replicates. Two-sided *t*-tests were used for accessing the phenotype difference between PH-1 and *Fgatg11* mutant. 'ns' means not significant. [Color figure can be viewed at wileyonlinelibrary.com]

identified after screening over a dozen hygromycin-resistant transformants. All of them had the same phenotypes described below although only data for one *Fgatg11* mutant were presented here. The *Fgatg11* mutants had no obvious defects in growth rate, plant infection, DON production, conidiation and conidia germination (Fig. 8A–F). Interestingly, although they were also normal in the formation of perithecia, asci, ascospores and ascospore germination (Fig. 8G–I), the *Fgatg11* mutants were defective in ascospore releasing (Fig. 8J). Whereas massive ascospores were forcibly discharged from mature perithecia in the wild type, only a small portion of the ascospores were ejected from *Fgatg11* perithecia (Fig. 8J). These results showed that *FgATG11* and selective autophagy play a stage-specific role during sexual reproduction in *F. graminearum*. Based on published RNA-seq data, *FgATG11* is constitutively expressed. Bioinformatics analysis also showed that *FgATG11* lacked stage-specific alternative splicing or transcription start/stop sites during sexual reproduction. Therefore, it is likely that stage-specific RNA editing may be related to or responsible for its functions during ascosporeogenesis.

Discussion

Autophagy is a dynamic and highly inducible degradation system that responds to environmental and physiological changes (Mizushima and Levine, 2010; Farre and Subramani, 2016) and involves various autophagy-related genes (Feng *et al.*, 2014). Investigation of the conservation and evolution of ATG genes in fungi is important for better understanding of the formation, evolution and regulation of autophagy. In this study, we systematically identified and analyzed the ATG genes in fungi with sequenced genomes. Our large-scale analysis revealed that only 20 ATG genes are highly conserved across the fungal kingdom, which is approximately half of those ATG genes characterized in the budding yeast, a model for studying autophagy. Interestingly, 14 ATG genes based on yeast homologues are unique to Saccharomycetes species and the other 7 are only present in Ascomycota. We also showed that important functional groups involved in the autophagosome formation contains conserved and non-conserved Atg components. In fact, many of the fungal species lacks the non-conserved components still possess

Fig. 9. Illustration of the loss and duplication of ATG genes in kingdom Fungi. The tree shows the evolutionary relationship of major groups of fungi. During the evolution of fungi, many ATG genes were gained but some were lost at different branching time. Independent losses and duplications of ATG genes in the fungal tree of life were labelled by the red and green patterns as marked respectively. [Color figure can be viewed at wileyonlinelibrary.com]



autophagy process. These non-conserved Atg proteins may be dispensable or replaceable in particular species, such as the absence of *ATG17* in the human pathogen *Cryptococcus neoformans*, which requires autophagy during infection (Hu *et al.*, 2008). Mammalian cells lack Atg17 ortholog but have a functional counterpart (Hara and Mizushima, 2009).

The ATG genes were completely or almost completely lost in Microsporidia and *Pneumocystis* that belong to different fungal lineages (Fig. 9). The massive losses of ATG genes are likely associated with their nutrient acquisition styles. As obligate intracellular pathogens, Microsporidia and *Pneumocystis* directly acquire nutrients from the host cell. The autophagy recycling system may be no longer required for these fungi to grow and reproduce within their hosts. These fungi have relative small genomes and lack many genes involved in primary metabolism and cellular signalling (Corradi, 2015; Ma *et al.*, 2016). Most likely, the ATG genes were lost independently in these fungi as parts of their genome reduction during the evolution. Differences among *Pneumocystis* species in the presence of a few specific ATG genes indicate that besides massive gene loss events occurred in their common ancestor, some of the ATG genes may be lost after their divergence. Similarly, some Microsporidia species differed from the majority of them in the distribution of ATG genes. Therefore, independent gene loss events likely occurred not only among different fungal lineages but also in a species-specific manner for at least some of these ATG genes in these obligate pathogens.

It is notable that massive losses of ATG genes are not a general feature of obligate or intracellular pathogens. The rust (basidiomycetes) and powdery mildew (ascomycetes) fungi are obligate plant pathogens that still have the core ATG genes but they are not intracellular pathogens. Some intracellular chytrid pathogens of animals such as the frog pathogen *Batrachochytrium dermatitidis* still have the core ATG genes but they are not obligate pathogens. Some chytrids, such as *Synchytridium endobioticum*, are obligate, intracellular plant pathogens but none of them has been sequenced. However, *Rozella allomyces* belonging to Cryptomycota is an obligate intracellular pathogen of other chytrids that still has most of the core ATG genes. The massive losses of ATG genes is also not a common feature related to infection of human or animal cells, because facultative pathogens such as *C. neoformans*, *Histoplasma capsulatum*, *Sporothrix schenckii* and *Blastomyces dermatitidis* still have most of the ATG genes (Casadevall, 2008). In comparison with other ATG genes subjected to massive losses, the retaining of *ATG15* in *Pneumocystis* and some Microsporidia species is also an interesting phenomenon. It is possible that the Atg15 lipase may have functions in other biological processes that are not directly related to autophagy. Interestingly, duplication of *ATG15* was observed in most Agaricomycetes species that tend to live in environments with limited lipid sources.

The third group of fungi with massive losses of ATG genes consists of three *Malassezia* species. Being smaller than 9-Mb, their genomes are among the smallest

genomes in the free-living basidiomycetes and filamentous ascomycetes (Xu *et al.*, 2007; Coelho *et al.*, 2013). Hence, the loss of *ATG* genes in *Malassezia* may be also associated with its reduction in genome size. However, unlike Microsporidia or *Pneumocystis*, *Malassezia* species are not obligate fungal pathogens although they have slow growth rate and lack sexual reproduction. *Malassezia* fungi can use lipids as the sole carbon source but fail to ferment sugars and have reduced numbers of glycosyl hydrolase and fatty acid synthase genes (Coelho *et al.*, 2013). With adaptation to animal skin environments with continuous lipid supplies, autophagy may be no longer essential for their growth and survivals in this niche. Strikingly, the nematode trapping fungus *D. stenobrocha* lacks *ATG1* and 8 other *ATG* genes, likely being defective in autophagy. However, autophagy is essential for nematode trapping in another Orbiliomycetes species, *A. oligospora* (Chen *et al.*, 2013), suggesting that losses of *ATG* genes likely occurred recently in *D. stenobrocha*. Nevertheless, unlike *D. stenobrocha*, *A. oligospora* forms extensive 3D adhesive networks to trap nematodes instead of constricting rings. In general, nematode-trapping fungi that form adhesive nets are more abundant in relatively poor soil and constricting ring-forming fungi favour environments rich in organic matters. Loss of *ATG* genes in *D. stenobrocha* may be related to its adaption to the organic compound-rich environments and its genome is also 11-Mb smaller than that of *A. oligospora* (Liu *et al.*, 2014). Taken together, all the massive loss of *ATG* genes observed in specific fungal lineages or groups are likely associated with their reduction in the genome size and adaptation to specific modes of nutrient acquirements.

Unlike massive losses of *ATG* genes that are observed in specific fungal groups as described above, our analysis showed that losses of specific or individual *ATG* genes have occurred independently throughout the fungal kingdom. Some of these species- or lineage-specific *ATG* gene loss events appeared to occur relatively recently because of their differences in distribution among closely related species. Even in Saccharomycetes, seven group II *ATG* genes were lost in 17–32 species. In higher fungi (filamentous ascomycetes and basidiomycetes), species- or lineage-specific (independent) losses were observed for several core and selective autophagy genes, including *ATG10*, *ATG16*, *ATG22* and *ATG33*. Some of these *ATG* gene loss events may be also related to their adaption to life or infection cycles, such as the loss of *ATG22* in Ustilagomycotina species. Many smut fungi have the free-living yeast form but obligate biotrophic hyphal form.

Besides gene losses, fungi may also gain specific *ATG* genes during evolution (Fig. 9). The basal fungal groups, such as Chytridiomycota and Mucoromycotina, have most of the group I highly conserved *ATG* genes. However, they lack all the *ATG* genes that are specific for

Saccharomycetes and likely are evolved after their divergence from the other ascomycetes. *ATG22*, a highly conserved core machinery *ATG* gene, is present in most of the Ascomycota and Basidiomycota species but absent in the basal fungal groups. Chytridiomycota and Mucoromycotina also lack *ATG10* and *ATG16*, two other core machinery *ATG* genes that are only conserved in Ascomycota. However, unlike *ATG22*, *ATG10* and *ATG16* are absent in Basidiomycetes. Lineage-specific gain of *ATG* genes also was observed for some selective autophagy specific genes, such as *ATG19* and *ATG34* in *Saccharomyces*. It is possible that the different conservation of these *ATG* gene are due to different *ATG* gene gained/evolved in the different stages of fungal evolution.

Lineage-specific duplications of certain *ATG* genes also were observed in different fungal groups. Interestingly, all the filamentous ascomycetes have a single copy of *ATG18* and *ATG20* but both of them are duplicated independently in the budding yeasts and fission yeasts. Whereas hyphae of filamentous ascomycetes have simple septa with septal pores that allow the free flow of mitochondria and other organelles between different compartments, the budding or fission yeasts are individual cells with rigid cell wall. Considering the functions of *Atg18* and *Atg20*, their duplications may contribute to the intracellular recycling of nutrients and organelles by differentially controlling the vesicle formation in autophagy and Cvt pathways in the budding and fission yeasts. Another interesting observation is that whereas duplication of *ATG15* occurred in Agariomycetes, *ATG22* was specifically duplicated in Ascomycetes. Although *Atg15* and *Atg22* differ in biological functions, both of them are involved in the final stages of autophagy. The differences in these gene duplication events suggest that different fungi may differ in nutrient recycling during autophagy.

Interestingly, unlike the other fungi, the dermatophytes in Arthrodermataceae possess two *ATG8* genes. To our knowledge, this is the first observation of a group of fungi with two *ATG8* paralogs of significant sequence divergence. The *ATG8* genes have evolved to three subfamilies in animals (Shpilka *et al.*, 2011) and also have been dramatically expanded in plants (Kellner *et al.*, 2016; Seo *et al.*, 2016). The three subfamilies of animal *Atg8s* differ in their subcellular localization, expression profiles and functions in the autophagy process (Shpilka *et al.*, 2011). *Dictyostelium discoideum* also has two *ATG8* genes that have different expression profiles and functions (Matthias *et al.*, 2016). In *T. rubrum*, the two *ATG8* genes also differed in the expression profiles, suggesting that they may also have distinct roles in the autophagy process. Because the natural habitats of the Arthrodermataceae dermatophytes are nutrient-limited, the two copies of *ATG8* may contribute to their adaptation the harsh environments. It is also possible that duplication of *ATG8* is

related to the differentiation of specific ornaments on their primitive fruiting bodies or the production of arthrospores on vegetative hyphae in Arthrodermataceae fungi.

Mucoromycotina species are the only other group of fungi that have duplication of *ATG8*. However, unlike Arthrodermataceae fungi, Mucoromycetes with two *ATG8*s also have many other *ATG* genes duplicated in the genome. Ancient whole-genome duplication may be responsible for the duplication of many genes in Mucoromycetes, including the genes involved in ergosterol synthesis (Ma *et al.*, 2009). The two *ATG8* paralogs in Mucoromycetes are highly similar to each other in amino acid sequences, which is different from Atg8a and Atg8b in Arthrodermataceae that have significant sequence divergence. It is likely that the two *ATG8* genes in Mucoromycetes have not undergone sequence and functional divergence yet due to their recent occurrence. Recent duplication of *ATG8* also was observed in three Agaricomycetes, *Pleurotus ostreatus*, *Sphaerobolus stellatus* and *Dacryopinax* species DJM-731 SS1 that have two *ATG8* genes encoding proteins of highly similar amino acid sequences. It will be interesting to determine the functions of *ATG8* duplication in these fungi. Because the expression level of *ATG8* is known to be related to the size of autophagosomes in other organisms (Xie *et al.*, 2008), it will be interesting to test whether two copies of *ATG8* will increase its overall expression and autophagosome size in these fungi.

Interestingly, approximately 90% of the *ATG* genes in *F. graminearum* and *N. crassa* are targeted by RNA editing during sexual reproduction. Since the A-to-I editing in fungi is generally adaptive (Wang *et al.*, 2016; Liu *et al.*, 2017), the editing of these *ATG* genes may be important for the autophagy function during the sexual reproduction. It is notable that although fungi have a single *ATG8* gene in general, different isoforms of Atg8 proteins can be generated by RNA editing during sexual reproduction. For example, both *F. graminearum* and *N. crassa* have two nonsynonymous editing sites in *ATG8* transcripts, which may lead to the synthesis of 4 isoforms of Atg8 proteins that may be slightly different in functions. These editing events in *ATG8* may be necessary for the fine-tuning of autophagy during sexual reproduction or provide protein heterozygosity beneficial for adaptation in the nature. *ATG13* is important for the initialization of both non-selective and selective autophagy. The Atg13 protein is targeted by the TORC1 kinase for phosphorylation. Inhibition of TORC1 leads to the hypophosphorylation of Atg13 and affects its interaction with Atg1 and Atg17 for the initiation of autophagy (Farre and Subramani, 2016). In *F. graminearum*, *N. crassa* and *N. tetrasperma*, the conserved editing event in *ATG13* results in the recoding of serine 536 to glycine, which may mimic the hypophosphorylation status of Atg13 proteins and affect autophagy

during sexual reproduction. In *F. graminearum*, deletion of *FgATG11*, the *ATG* gene with the most editing sites in *F. graminearum*, had no obvious effect on growth, conidiation, plant infection and formation of perithecia or ascospores but specifically affected forcibly discharge of ascospores, suggesting a relationship between stage-specific function and RNA editing of *FgATG11* during sexual reproduction. Deletion of *PUK1* and *AMD1*, two other genes with sexual stage-specific RNA editing, also affected ascospore release in *F. graminearum* (Liu *et al.*, 2016; Cao *et al.*, 2017). Overall, our results indicated that RNA editing may affect the expression or sequence of Atg proteins and play a role in fine tuning the autophagy process during ascospore formation and release. Therefore, it will be important to functionally characterize the functions of *ATG* genes and their editing events during sexual reproduction in *F. graminearum* and *N. crassa*.

Methods

Reference ATG genes from yeasts and filamentous fungi

The sequences of 36 *ATG* genes of *S. cerevisiae* (Supporting Information Table S1) were retrieved from SGD (yeastgenome.org). The sequences of Atg25 from the methylotrophic yeast *Hansenula polymorpha* and Atg28, Atg30, Atg35 and Atg37 from the methylotrophic yeast *Pichia pastoris* (Supporting Information Table S1) were based on earlier publications (Monastyrska *et al.*, 2005; Meijer *et al.*, 2007). The *M. oryzae* and *F. oxysporum* *ATG* genes (Kershaw and Talbot, 2009; Corral-Ramos *et al.*, 2015) were manually verified by both bidirectional blast and OrthoMCL (Li *et al.*, 2003) methods. The *ATG8* and *ATG15* genes of *F. graminearum* were obtained from earlier publications (Nguyen *et al.*, 2011; Josefsen *et al.*, 2012). All the other *ATG* genes identified in *F. graminearum* were manually annotated.

Genome sequences and RNA-seq data of different fungi

Most of the genome sequences and annotations used in this study were retrieved from the Ensembl Fungi genome database (fungi.ensembl.org/index.html) release 34. The genome sequences of *Phycomyces blakesleeianus* (Corrochano *et al.*, 2016), *Taphrina deformans* (Cisse *et al.*, 2013), *Gonapodya prolifera* (Chang *et al.*, 2015) and *M. globosa* (Xu *et al.*, 2007) were downloaded from the DOE Joint Genome Institute. The *P. jirovecii*, *P. carinii* (Ma *et al.*, 2016), *N. antheraeae* (Pan *et al.*, 2013) and *Hamiltosporidium tvaerminnensis* (Corradi *et al.*, 2009) genomes were downloaded from NCBI genome database. The genomes of *Mortierella verticillata*, *Punctularia strigosozonata*, *Aspergillus ustus*, *Talaromyces cellulolyticus*, *Fomitiporia mediterranea*, *Trametes versicolor*, *Stereum*

hirsutum, *Plicaturopsis crispa*, *Coniophora puteana*, *Tremella mesenterica* and *Rhodotorula glutinis* were not well assembled or annotated. For *Colletotrichum higginsianum*, *Verticillium longisporum* and *Allomyces macrogynus*, their genome sequences appeared to be derived from heterokaryotic or diploid hyphae. These fungal species were excluded from this analysis to avoid false negative or false positive results with ATG gene annotations.

The raw reads of *T. rubrum* RNA-seq data (Xu *et al.*, 2015) were mapped to its genome by hisat2 (Kim *et al.*, 2015). The bam file was generated by SAMtools (Li *et al.*, 2009) and sashimi plots were generated with IGV (Thorvaldsdottir *et al.*, 2013). The 3D structures of TrAtg8s were predicted with the I-TASSER server (Yang *et al.*, 2015) and visualized with PyMol (pymol.org).

Identification of fungal ATG genes

To systematically identify fungal ATG genes, the in-house Perl scripts (available at github.com/wangqinhu/fungi_atg) were developed and used to retrieve and parse homology information from release 34 of Ensembl Fungi. To ensure that all the orthologs were correctly identified, four previously reported and manually verified seed sequences were applied for most ATG genes. For the genomes that are not included in the genome database of Ensembl Fungi, a bidirectional blastp search was performed to define the orthologs. For species with most of the ATG gene but lacking an ATG8 ortholog, a tblastn search was performed against the genome sequence to identify potential mis-annotation. For ATG1, ATG10 and ATG26, manual annotation also was applied to correct errors introduced by automated annotation. For *Dichomitus squalens*, *Glarea lozoyensis*, *Gymnopus luxurians*, *Hydnomerulius pinastri*, *Laccaria bicolor*, *Paxillus involutus*, *Postia placenta*, *Pseudozyma antarctica*, *Pseudozyma aphidis* and *Sphaerobolus stellatus*, the annotation of ATG genes were corrected with the data available from the DOE Joint Genome Institute by blastp or tblastn searches.

Phylogenetic analysis

The sequence alignments were generated with M-Coffee (Wallace *et al.*, 2006) and adjusted with trimAl (Capella-Gutiérrez *et al.*, 2009). The phylogenetic relationships were inferred with the maximum likelihood method as developed in PhyML 3.1 (Guindon and Gascuel, 2003). The trees were visualized by FigTree (tree.bio.ed.ac.uk/software/figtree).

Editing sites in ATG genes of *F. graminearum* and *N. crassa*

The A-to-I editing data of *F. graminearum* and *N. crassa* were generated in previous studies (Liu *et al.*, 2016,

2017). The editing sites in ATG genes were plotted with costumed R script. Conserved A-to-I editing sites in *F. graminearum*, *N. crassa* and *N. tetrasperma* were also obtained from previous report (Liu *et al.*, 2016).

Generation and characterization of the ATG genes mutant in *F. graminearum*

To functionally characterize the two editing sites in *FgATG8*, we generated mutant alleles of *FgATG8* carrying the T93G, A94G, T120G, or A121G mutation by overlapping PCR. The resulting *FgATG8*^{G93}, *FgATG8*^{G94}, *FgATG8*^{G120} and *FgATG8*^{G121} constructs were verified by sequencing analysis and transformed into the *Fgatg8* mutant generated by the split-marker approach (Catlett *et al.*, 2002). Transformants of *Fgatg8* expressing these mutant alleles were confirmed by PCR analysis and assayed for defects in sexual reproduction as described (Josefsen *et al.*, 2012).

To generate the *FgATG11* gene replacement mutant, its upstream and downstream flanking sequences were amplified and fused with the N-terminal portion and C-terminal portion of the hygromycin phosphotransferase (*hph*) cassette (Catlett *et al.*, 2002), respectively, by overlapping PCR. The resulting PCR products were transformed into the wild-type strain PH-1 as described (Zhou *et al.*, 2011). The resulting hygromycin-resistant transformants were screened by PCR analysis (Supporting Information Fig. S3). Growth rate, infection assays with flowering wheat heads, disease index, conidiation and conidium germination were assayed as previously described (Wang *et al.*, 2011; Cao *et al.*, 2016; Wang *et al.*, 2018). Perithecium formation, ascus development, ascospore germination and ascospore releasing were assayed on carrot agar plates as described (Cavinder *et al.*, 2012; Luo *et al.*, 2014; Yin *et al.*, 2018).

Acknowledgements

We thank Dr. Tingting Li for improving the manuscript and fruitful discussion. This work was supported by grants from National Key R&D Program of China (2017YFD0200602-2), National Natural Science Foundation of China (31701747 and 31601587), China Postdoctoral Science Foundation (2015M580884), National Excellent Doctoral Dissertations (201471), and US Wheat and Barley Scab Initiative.

References

- Ashbee, H. R., and Evans, E. G. (2002) Immunology of diseases associated with *Malassezia* species. *Clin Microbiol Rev* **15**: 21–57.
- Avila-Ospina, L., Moison, M., Yoshimoto, K., and Masclaux-Daubresse, C. (2014) Autophagy, plant senescence, and nutrient recycling. *J Exp Bot* **65**: 3799–3811.

- Barth, H., Meiling-Wesse, K., Epple, U. D., and Thumm, M. (2001) Autophagy and the cytoplasm to vacuole targeting pathway both require Aut10p. *FEBS Lett* **508**: 23–28.
- Bassham, D. C., Laporte, M., Marty, F., Moriyasu, Y., Ohsumi, Y., Olsen, L. J., and Yoshimoto, K. (2006) Autophagy in development and stress responses of plants. *Autophagy* **2**: 2–11.
- Cao, S., Zhang, S., Hao, C., Liu, H., Xu, J. R., and Jin, Q. (2016) FgSsn3 kinase, a component of the mediator complex, is important for sexual reproduction and pathogenesis in *Fusarium graminearum*. *Sci Rep* **6**: 22333.
- Cao, S., He, Y., Hao, C., Xu, Y., Zhang, H., Wang, C., et al. (2017) RNA editing of the *AMD1* gene is important for ascus maturation and ascospore discharge in *Fusarium graminearum*. *Sci Rep* **7**: 4617.
- Capella-Gutierrez, S., Silla-Martinez, J. M., and Gabaldon, T. (2009) trimAl: a tool for automated alignment trimming in large-scale phylogenetic analyses. *Bioinformatics* **25**: 1972–1973.
- Casadevall, A. (2008) Evolution of intracellular pathogens. *Annu Rev Microbiol* **62**: 19–33.
- Cattlett, N. L., Lee, B.-N., Yoder, O. C., and Turgeon, B. G. (2002) Split-marker recombination for efficient targeted deletion of fungal genes. *Fungal Genet News* **50**: 9–11.
- Cavinder, B., Sikhakolli, U., Fellows, K. M., and Trail, F. (2012) Sexual development and ascospore discharge in *Fusarium graminearum*. *J Vis Exp* 3895. doi:10.3791/3895
- Chang, Y., Wang, S., Sekimoto, S., Aerts, A. L., Choi, C., Clum, A., et al. (2015) Phylogenomic analyses indicate that early fungi evolved digesting cell walls of algal ancestors of land plants. *Genome Biol Evol* **7**: 1590–1601.
- Chen, Y. L., Gao, Y., Zhang, K. Q., and Zou, C. G. (2013) Autophagy is required for trap formation in the nematode-trapping fungus *Arthrobotrys oligospora*. *Environ Microbiol Rep* **5**: 511–517.
- Cisse, O. H., Almeida, J. M., Fonseca, A., Kumar, A. A., Salojarvi, J., Overmyer, K., et al. (2013) Genome sequencing of the plant pathogen *Taphrina deformans*, the causal agent of peach leaf curl. *MBio* **4**: e00055-00013.
- Coelho, M. A., Sampaio, J. P., and Goncalves, P. (2013) Living and thriving on the skin: *Malassezia* genomes tell the story. *MBio* **4**: e00117–e00113.
- Corradi, N. (2015) Microsporidia: eukaryotic intracellular parasites shaped by gene loss and horizontal gene transfers. *Annu Rev Microbiol* **69**: 167–183.
- Corradi, N., Haag, K. L., Pombert, J. F., Ebert, D., and Keeling, P. J. (2009) Draft genome sequence of the *Daphnia* pathogen *Octospora bayeri*: insights into the gene content of a large microsporidian genome and a model for host-parasite interactions. *Genome Biol* **10**: R106.
- Corral-Ramos, C., Roca, M. G., Di Pietro, A., Roncero, M. I., and Ruiz-Roldan, C. (2015) Autophagy contributes to regulation of nuclear dynamics during vegetative growth and hyphal fusion in *Fusarium oxysporum*. *Autophagy* **11**: 131–144.
- Corrochano, L. M., Kuo, A., Marcet-Houben, M., Polaino, S., Salamov, A., Villalobos-Escobedo, J. M., et al. (2016) Expansion of signal transduction pathways in fungi by extensive genome duplication. *Curr Biol* **26**: 1577–1584.
- Deng, Y. Z., Qu, Z., He, Y., and Naqvi, N. I. (2012) Sorting nexin Snx41 is essential for conidiation and mediates glutathione-based antioxidant defense during invasive growth in *Magnaporthe oryzae*. *Autophagy* **8**: 1058–1070.
- Deng, Y., Qu, Z., and Naqvi, N. I. (2013) The role of Snx41-based pexophagy in *Magnaporthe* development. *PLoS One* **8**: e79128.
- Dolatabadi, S., Scherlach, K., Figge, M., Hertweck, C., Dijksterhuis, J., Menken, S. B., and de Hoog, G. S. (2016) Food preparation with mucoralean fungi: a potential bio-safety issue? *Fungal Biol* **120**: 393–401.
- Duan, Z., Chen, Y., Huang, W., Shang, Y., Chen, P., and Wang, C. (2013) Linkage of autophagy to fungal development, lipid storage and virulence in *Metarhizium robertsii*. *Autophagy* **9**: 538–549.
- Eskelinen, E. L., and Saftig, P. (2009) Autophagy: a lysosomal degradation pathway with a central role in health and disease. *Biochim Biophys Acta* **1793**: 664–673.
- Farre, J. C., and Subramani, S. (2016) Mechanistic insights into selective autophagy pathways: lessons from yeast. *Nat Rev Mol Cell Biol* **17**: 537–552.
- Feng, Y., He, D., Yao, Z., and Klionsky, D. J. (2014) The machinery of macroautophagy. *Cell Res* **24**: 24–41.
- Geng, J., and Klionsky, D. J. (2008) The Atg8 and Atg12 ubiquitin-like conjugation systems in macroautophagy. 'Protein modifications: beyond the usual suspects' review series. *EMBO Rep* **9**: 859–864.
- Guindon, S., and Gascuel, O. (2003) A simple, fast, and accurate algorithm to estimate large phylogenies by maximum likelihood. *Syst Biol* **52**: 696–704.
- Haag, K. L., James, T. Y., Pombert, J. F., Larsson, R., Schaer, T. M., Refardt, D., and Ebert, D. (2014) Evolution of a morphological novelty occurred before genome compaction in a lineage of extreme parasites. *Proc Natl Acad Sci U S A* **111**: 15480–15485.
- Hara, T., and Mizushima, N. (2009) Role of ULK-FIP200 complex in mammalian autophagy: FIP200, a counterpart of yeast Atg17? *Autophagy* **5**: 85–87.
- Hetteima, E. H., Lewis, M. J., Black, M. W., and Pelham, H. R. (2003) Retromer and the sorting nexins Snx4/41/42 mediate distinct retrieval pathways from yeast endosomes. *EMBO J* **22**: 548–557.
- Hu, G., Hacham, M., Waterman, S. R., Panepinto, J., Shin, S., Liu, X., et al. (2008) PI3K signaling of autophagy is required for starvation tolerance and virulence of *Cryptococcus neoformans*. *J Clin Invest* **118**: 1186–1197.
- Josefsen, L., Droce, A., Sondergaard, T. E., Sorensen, J. L., Bormann, J., Schafer, W., et al. (2012) Autophagy provides nutrients for nonassimilating fungal structures and is necessary for plant colonization but not for infection in the necrotrophic plant pathogen *Fusarium graminearum*. *Autophagy* **8**: 326–337.
- Kanayama, M., and Shinohara, M. L. (2016) Roles of autophagy and autophagy-related proteins in antifungal immunity. *Front Immunol* **7**: 47.
- Keeling, P. (2009) Five questions about Microsporidia. *PLoS Pathog* **5**: e1000489.
- Kellner, R., De la Concepcion, J. C., Maqbool, A., Kamoun, S., and Dagdas, Y. F. (2016) ATG8 expansion: a driver of selective autophagy diversification? *Trends Plant Sci* **22**: 204–214.

- Kershaw, M. J., and Talbot, N. J. (2009) Genome-wide functional analysis reveals that infection-associated fungal autophagy is necessary for rice blast disease. *Proc Natl Acad Sci U S A* **106**: 15967–15972.
- Kim, D., Langmead, B., and Salzberg, S. L. (2015) HISAT: a fast spliced aligner with low memory requirements. *Nat Methods* **12**: 357–360.
- Krick, R., Henke, S., Tolstrup, J., and Thumm, M. (2008) Dissecting the localization and function of Atg18, Atg21 and Ygr223c. *Autophagy* **4**: 896–910.
- Levine, B., Mizushima, N., and Virgin, H. W. (2011) Autophagy in immunity and inflammation. *Nature* **469**: 323–335.
- Li, L., Stoeckert, C. J., Jr., and Roos, D. S. (2003) OrthoMCL: identification of ortholog groups for eukaryotic genomes. *Genome Res* **13**: 2178–2189.
- Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., et al. (2009) The sequence alignment/map format and SAMtools. *Bioinformatics* **25**: 2078–2079.
- Liu, Y., Schiff, M., Czymmek, K., Tallozy, Z., Levine, B., and Dinesh-Kumar, S. P. (2005) Autophagy regulates programmed cell death during the plant innate immune response. *Cell* **121**: 567–577.
- Liu, X. H., Gao, H. M., Xu, F., Lu, J. P., Devenish, R. J., and Lin, F. C. (2012) Autophagy vitalizes the pathogenicity of pathogenic fungi. *Autophagy* **8**: 1415–1425.
- Liu, K., Zhang, W., Lai, Y., Xiang, M., Wang, X., Zhang, X., and Liu, X. (2014) *Drechlerella stenobrocha* genome illustrates the mechanism of constricting rings and the origin of nematode predation in fungi. *BMC Genomics* **15**: 114.
- Liu, H., Wang, Q., He, Y., Chen, L., Hao, C., Jiang, C., et al. (2016) Genome-wide A-to-I RNA editing in fungi independent of ADAR enzymes. *Genome Res* **26**: 499–509.
- Liu, H., Li, Y., Chen, D., Qi, Z., Wang, Q., Wang, J., et al. (2017) A-to-I RNA editing is developmentally regulated and generally adaptive for sexual reproduction in *Neurospora crassa*. *Proc Natl Acad Sci U S A* **114**: E7756–E7765.
- Luo, Y., Zhang, H., Qi, L., Zhang, S., Zhou, X., Zhang, Y., and Xu, J. R. (2014) FgKin1 kinase localizes to the septal pore and plays a role in hyphal growth, ascospore germination, pathogenesis, and localization of Tub1 beta-tubulins in *Fusarium graminearum*. *New Phytol* **204**: 943–954.
- Lv, W., Wang, C., Yang, N., Que, Y., Talbot, N. J., and Wang, Z. (2017) Genome-wide functional analysis reveals that autophagy is necessary for growth, sporulation, deoxynivalenol production and virulence in *Fusarium graminearum*. *Sci Rep* **7**: 11062.
- Lynch-Day, M. A., and Klionsky, D. J. (2010) The Cvt pathway as a model for selective autophagy. *FEBS Lett* **584**: 1359–1366.
- Ma, L. J., Ibrahim, A. S., Skory, C., Grabherr, M. G., Burger, G., Butler, M., et al. (2009) Genomic analysis of the basal lineage fungus *Rhizopus oryzae* reveals a whole-genome duplication. *PLoS Genet* **5**: e1000549.
- Ma, L., Chen, Z., da Huang, W., Kutty, G., Ishihara, M., Wang, H., et al. (2016) Genome analysis of three pneumocystis species reveals adaptation mechanisms to life exclusively in mammalian hosts. *Nat Commun* **7**: 10740.
- Matthias, J., Messling, S., and Eichinger, L. (2016) The two *Dictyostelium* autophagy eight proteins, ATG8a and ATG8b, associate with the autophagosome in succession. *Eur J Cell Biol* **95**: 15–25.
- Meijer, W. H., van der Klei, I. J., Veenhuis, M., and Kiel, J. A. (2007) ATG genes involved in non-selective autophagy are conserved from yeast to man, but the selective Cvt and pexophagy pathways also require organism-specific genes. *Autophagy* **3**: 106–116.
- Michaeli, S., Galili, G., Genschik, P., Fernie, A. R., and Avin-Wittenberg, T. (2016) Autophagy in plants – what's new on the menu? *Trends Plant Sci* **21**: 134–144.
- Mizushima, N. (2007) Autophagy: process and function. *Genes Dev* **21**: 2861–2873.
- Mizushima, N., and Levine, B. (2010) Autophagy in mammalian development and differentiation. *Nat Cell Biol* **12**: 823–830.
- Mizushima, N., Levine, B., Cuervo, A. M., and Klionsky, D. J. (2008) Autophagy fights disease through cellular self-digestion. *Nature* **451**: 1069–1075.
- Monastyrska, I., Kiel, J. A., Krikken, A. M., Komduur, J. A., Veenhuis, M., and van der Klei, I. J. (2005) The *Hansenula polymorpha* ATG25 gene encodes a novel coiled-coil protein that is required for macropexophagy. *Autophagy* **1**: 92–100.
- Nadal, M., and Gold, S. E. (2010) The autophagy genes ATG8 and ATG1 affect morphogenesis and pathogenicity in *Ustilago maydis*. *Mol Plant Pathol* **11**: 463–478.
- Nair, U., Cao, Y., Xie, Z., and Klionsky, D. J. (2010) Roles of the lipid-binding motifs of Atg18 and Atg21 in the cytoplasm to vacuole targeting pathway and autophagy. *J Biol Chem* **285**: 11476–11488.
- Nguyen, L. N., Bormann, J., Le, G. T., Starkel, C., Olsson, S., Nosanchuk, J. D., et al. (2011) Autophagy-related lipase FgATG15 of *Fusarium graminearum* is important for lipid turnover and plant infection. *Fungal Genet Biol* **48**: 217–224.
- Ohsumi, Y. (2001) Molecular dissection of autophagy: two ubiquitin-like systems. *Nat Rev Mol Cell Biol* **2**: 211–216.
- Pan, G., Xu, J., Li, T., Xia, Q., Liu, S. L., Zhang, G., et al. (2013) Comparative genomics of parasitic silkworm microsporidia reveal an association between genome expansion and host adaptation. *BMC Genomics* **14**: 186.
- Pollack, J. K., Harris, S. D., and Marten, M. R. (2009) Autophagy in filamentous fungi. *Fungal Genet Biol* **46**: 1–8.
- Roden, M. M., Zaoutis, T. E., Buchanan, W. L., Knudsen, T. A., Sarkisova, T. A., Schaufele, R. L., et al. (2005) Epidemiology and outcome of zygomycosis: a review of 929 reported cases. *Clin Infect Dis* **41**: 634–653.
- Schwartz, V. U., Winter, S., Shelest, E., Marcet-Houben, M., Horn, F., Wehner, S., et al. (2014) Gene expansion shapes genome architecture in the human pathogen *Lichtheimia corymbifera*: an evolutionary genomics analysis in the ancient terrestrial mucorales (Mucoromycotina). *PLoS Genet* **10**: e1004496.
- Seo, E., Woo, J., Park, E., Bertolani, S. J., Siegel, J. B., Choi, D., and Dinesh-Kumar, S. P. (2016) Comparative analyses of ubiquitin-like ATG8 and cysteine protease ATG4 autophagy genes in the plant lineage and cross-kingdom processing of ATG8 by ATG4. *Autophagy* **12**: 2054–2068.

- Shpilka, T., Weidberg, H., Pietrokovski, S., and Elazar, Z. (2011) Atg8: an autophagy-related ubiquitin-like protein family. *Genome Biol* **12**: 226.
- Stromhaug, P. E., Reggiori, F., Guan, J., Wang, C. W., and Klionsky, D. J. (2004) Atg21 is a phosphoinositide binding protein required for efficient lipidation and localization of Atg8 during uptake of aminopeptidase I by selective autophagy. *Mol Biol Cell* **15**: 3553–3566.
- Suzuki, H., Osawa, T., Fujioka, Y., and Noda, N. N. (2016) Structural biology of the core autophagy machinery. *Curr Opin Struct Biol* **43**: 10–17.
- Thorvaldsdottir, H., Robinson, J. T., and Mesirov, J. P. (2013) Integrative genomics viewer (IGV): high-performance genomics data visualization and exploration. *Brief Bioinform* **14**: 178–192.
- Tsukada, M., and Ohsumi, Y. (1993) Isolation and characterization of autophagy-defective mutants of *Saccharomyces cerevisiae*. *FEBS Lett* **333**: 169–174.
- Voigt, O., and Poggeler, S. (2013) Autophagy genes *Smatg8* and *Smatg4* are required for fruiting-body development, vegetative growth and ascospore germination in the filamentous ascomycete *Sordaria macrospora*. *Autophagy* **9**: 33–49.
- Wallace, I. M., O'Sullivan, O., Higgins, D. G., and Notredame, C. (2006) M-coffee: combining multiple sequence alignment methods with T-coffee. *Nucleic Acids Res* **34**: 1692–1699.
- Wang, C., Zhang, S., Hou, R., Zhao, Z., Zheng, Q., Xu, Q., et al. (2011) Functional analysis of the kinome of the wheat scab fungus *Fusarium graminearum*. *PLoS Pathog* **7**: e1002460.
- Wang, Q., Jiang, C., Liu, H., and Xu, J.-R. (2016) ADAR-independent A-to-I RNA editing is generally adaptive for sexual reproduction in fungi. *BioRxiv* : 059725. doi:10.1101/059725
- Wang, Q., Chen, D., Wu, M., Zhu, J., Jiang, C., Xu, J. R., and Liu, H. (2018) MFS transporters and GABA metabolism are involved in the self-defense against DON in *Fusarium graminearum*. *Front Plant Sci* **9**: 438.
- Xie, Z., Nair, U., and Klionsky, D. J. (2008) Atg8 controls phagophore expansion during autophagosome formation. *Mol Biol Cell* **19**: 3290–3298.
- Xu, J., Saunders, C. W., Hu, P., Grant, R. A., Boekhout, T., Kuramae, E. E., et al. (2007) Dandruff-associated *Malassezia* genomes reveal convergent and divergent virulence traits shared with plant and human fungal pathogens. *Proc Natl Acad Sci U S A* **104**: 18730–18735.
- Xu, X., Liu, T., Ren, X., Liu, B., Yang, J., Chen, L., et al. (2015) Proteogenomic analysis of *Trichophyton rubrum* aided by RNA sequencing. *J Proteome Res* **14**: 2207–2218.
- Yang, J., Yan, R., Roy, A., Xu, D., Poisson, J., and Zhang, Y. (2015) The I-TASSER suite: protein structure and function prediction. *Nat Methods* **12**: 7–8.
- Yao, Z., Delorme-Axford, E., Backues, S. K., and Klionsky, D. J. (2015) Atg41/lcy2 regulates autophagosome formation. *Autophagy* **11**: 2288–2299.
- Yin, T., Zhang, Q., Wang, J., Liu, H., Wang, C., Xu, J. R., and Jiang, C. (2018) The cyclase-associated protein FgCap1 has both protein kinase A-dependent and -independent functions during deoxynivalenol production and plant infection in *Fusarium graminearum*. *Mol Plant Pathol* **19**: 552–563.
- Yoshimori, T., and Noda, T. (2008) Toward unraveling membrane biogenesis in mammalian autophagy. *Curr Opin Cell Biol* **20**: 401–407.
- Zhao, D., Liu, X. M., Yu, Z. Q., Sun, L. L., Xiong, X., Dong, M. Q., and Du, L. L. (2016) Atg20- and Atg24-family proteins promote organelle autophagy in fission yeast. *J Cell Sci* **129**: 4289–4304.
- Zhou, X., Li, G., and Xu, J.-R. (2011). Efficient approaches for generating GFP fusion and epitope-tagging constructs in filamentous fungi. In *Fungal Genomics: Methods and Protocols*, Xu, J.-R., and Bluhm, B. H. (eds). Totowa, NJ, USA: Humana Press, pp. 199–212.

Supporting Information

Additional Supporting Information may be found in the online version of this article at the publisher's web-site:

Fig. S1. The conservation of ATG genes in Saccharomycetes.

The dendrogram on the left shows the phylogenetic relationship of the marked Saccharomycetes species. The copy numbers of individual ATG genes (highly conserved, red; conserved, green; specific, yellow) in each species vary from 0 to 3. The ATG genes from *Fusarium graminearum*, *Magnaporthe oryzae* and *Neurospora crassa* are included as the outliers for comparison.

Fig. S2. Mutations affecting the editing events at A94 and A121 in FgATG8.

(A) Mating cultures of the wild type strain PH-1, *Fgatg8* deletion mutant, and transformants of the *Fgatg8* mutant expressing the non-editable *FgATG8^{G93}* and *FgATG8^{C120}* alleles or edited *FgATG8^{G94}* and *FgATG8^{G121}* alleles were assayed for perithecia formation at 8 days post-fertilization. (B) Asci and ascospores of PH-1 and transformants of the *Fgatg8* mutant expressing *FgATG8^{G93}*, *FgATG8^{C120}*, *FgATG8^{G94}*, and *FgATG8^{G121}* alleles. Bar = 20 μm.

Fig. S3. Diagram of the FgATG11 gene and its gene replacement construct.

Primer pairs F1/R2 and F3/R4 were used to amplify the upstream and downstream flanking sequences of *FgATG11* respectively. The N- and C-portions of the hygromycin phosphotransferase (hph) cassette were amplified with primer pairs HYG-F/HY-R and YG-F/HYG-R. The overlapping PCR products amplified by primer pairs F1/HY-R and YG-F/R4 were co-transformed into the wild-type strain PH-1. Hygromycin-resistant mutants were screened with primer pairs H852/H850, F5/R6, F7/H855-R and H856-F/R8.

Table S1. ATG genes in yeasts.

Table S2. The number of ATG genes in fungi.

Table S3. The abbreviations of species used for the phylogenetic analysis.